



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$Q_k m + n Q_{k-1} = P_k \dots (1),$$

$$P_k m + n P_{k-1} = Q_k A \dots (2).$$

Solving for  $n$  in (2) and making use of the relation  $P_k Q_{k-1} - Q_k P_{k-1} = (-1)^k$  we get,  $P_k^2 - A Q_k^2 = (-1)^k n$ .

Also solved by *G. B. M. ZERR*.

143. Proposed by *JOHN M. COLAW*, A. M., Monterey, Va.

Solve  $x + y + z + a = a \dots (1)$ .

$$x^2 + y^2 + z^2 + u^2 = b \dots (2).$$

$$x^3 + y^3 + z^3 + u^3 = c \dots (3).$$

$$x^4 + y^4 + z^4 + u^4 = d \dots (4).$$

Solution by *MARCUS BAKER*, Washington, D. C.

Let  $u$ ,  $x$ ,  $y$  and  $z$  be the roots of the equation

$$X^4 - AX^3 + BX^2 - CX + D = (X - u)(X - x)(X - y)(X - z).$$

For  $X$  write  $1/V$ , whence

$$1 - AV + BV^2 - CV^3 + DV^4 = (1 - uV)(1 - xV)(1 - yV)(1 - zV);$$

whence  $\log(1 - AV + BV^2 - CV^3 + DV^4)$

$$\begin{aligned} &= \log(1 - uV) + \log(1 - xV) + \log(1 - yV) + \log(1 - zV), \\ &= -uV - \frac{1}{2}u^2V^2 - \frac{1}{3}u^3V^3 - \frac{1}{4}u^4V^4 - \dots \\ &\quad -xV - \frac{1}{2}x^2V^2 - \frac{1}{3}x^3V^3 - \frac{1}{4}x^4V^4 - \dots \\ &\quad -yV - \frac{1}{2}y^2V^2 - \frac{1}{3}y^3V^3 - \frac{1}{4}y^4V^4 - \dots \\ &\quad -zV - \frac{1}{2}z^2V^2 - \frac{1}{3}z^3V^3 - \frac{1}{4}z^4V^4 - \dots \\ &= -aV - \frac{1}{2}bV^2 - \frac{1}{3}cV^3 - \frac{1}{4}dV^4 - \dots \end{aligned}$$

whence  $1 - AV + BV^2 - CV^3 + DV^4 = e^{-(aV + \frac{1}{2}bV^2 + \frac{1}{3}cV^3 + \frac{1}{4}dV^4 + \dots)}$

Developing the second member of this equation, remembering that

$$e^{-x} = 1 - x - \frac{x^2}{2!} - \frac{x^3}{3!} - \frac{x^4}{4!} \dots$$

we have

$$1 - AV + BV^2 - CV^3 + DV^4 = 1 - a \left\{ \begin{array}{l} V - \frac{1}{2}b \\ + \frac{1}{2}a^2 \end{array} \right\} V^3 - \frac{1}{3}c \left\{ \begin{array}{l} V - \frac{1}{2}d \\ + \frac{1}{2}ab \\ - \frac{1}{6}a^3 \end{array} \right\} V^4 - \dots$$

Equating like coefficients, we have

$$A = a; \quad B = \frac{1}{2}(a^2 - b); \quad C = \frac{1}{6}(a^3 - 3ab + 2c); \quad D = \frac{1}{24}[a^4 - 6a^2b + (3b^2 + 8ac) - 6d].$$

Therefore  $u$ ,  $x$ ,  $y$  and  $z$  are the roots of the equations

$$X^4 - aX^3 + \frac{1}{2}(a^2 - b)X^2 - \frac{1}{6}(a^3 - 3ab + 2c)X + \frac{1}{24}(a^4 - 6a^2b[8ac + 3b^2] - 6d) = 0.$$

The *method* here used I obtained many years ago from my old friend James Main of the Coast and Geodetic Survey who died in Washington November 23, 1894, aged 84 years. This method is *general* applying to a set of  $n$  equations containing  $n$  unknown quantities.

If we have

$$\begin{aligned} x + y + z + \dots &= a \\ x^2 + y^2 + z^2 + \dots &= b \\ x^3 + y^3 + z^3 + \dots &= c \\ \cdot &\quad \cdot \quad \cdot \quad \cdot \\ x^n + y^n + z^n + \dots &= k \end{aligned}$$

then are  $x, y, z$ , etc., the roots of

$$X^n - AX^{n-1} + BX^{n-2} - CX^{n-3} + \dots \pm K = 0,$$

where

$$A = a$$

$$B = (1/2!) [a^2 - b]$$

$$C = (1/3!) [a^3 - 3ab + 2c]$$

$$D = (1/4!) [a^4 - 6a^2b + (8ac + 3b^2) - 6d]$$

$$E = (1/5!) [a^5 - 10a^3b + 5a(4ac + 3b^2) - 10(3ad + 2bc) + 24c]$$

$$F = (1/6!) [a^6 - 15a^4b + 5a^2(8ac + 9b^2) - 15(6a^2d + 8abc + b^3) + 2(72ae + 45bd + 20c^2) - 120f]$$

$$G = (1/7!) [a^7 - 21a^5b + 35a^3(2ac + 3b^2) - 105a(2a^2d + 3bc + b^3 + b^2c) + 14(24a^2e + 20ac^2 + 15b^2c + 45abd) - 84(10af + 6be + 5cd) + 720g]$$

Solved in an excellent manner by G. B. M. ZERR.

144. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy, Defiance College, Defiance, Ohio.

Show that the number of ways in which 15 different problems may be distributed among 5 students so that each student shall have three of them, is  $N = (5.3)!/(3!)^5$ .

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa., and M. E. GRABER, Heidelberg University, Tiffin, Ohio.

It is demonstrated that the number of ways  $a + b + c + \dots$  can be divided into groups containing  $a$  things in one group,  $b$  in another, etc., is

$$\frac{[a + b + c + \dots]!}{[a!][b!][c!]\dots}$$